



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

"Napier based his explanations upon two considerations: (1) the geometrico-mechanical concept of flowing points, (2) the relations which exist between arithmetic and geometric series. . . . Napier lets the point g move along the definite line TS with a diminishing velocity such that its velocity at T is to that at d , as the distance TS is to the distance dS . At the same time Napier lets a point a move along the line bc (which is of indefinite length) with a uniform velocity which is the same as the initial velocity of the point g . If the two points start to move at the same moment, and if g is at d when a is at c , then the length bc is defined as the logarithm of dS . Napier constructed tables for trigonometric computation. With that end in view he lets TS stand for the *radius*, assigning to it the value 10^7 , while dS stands for a given *sine*. At that time trigonometric functions were not thought of strictly as *ratios*."

Then I give Napier's kinematical definition *in his own words*; it is the very definition in the *Constructio*, § 26, which Professor Carslaw says in the above communication that I ignore. What more could I do to emphasize this kinematical definition?

As an easy deduction from this kinematical definition is the one-one correspondence between arithmetic progressions and geometric progressions used by Napier in his computations. The part of Professor Carslaw's criticism which I consider valid is that $\text{Nap. log } (10^7 - 1) = 1.00000005$ exhibits "Table I" in Napier's *Constructio* somewhat more closely than does $\text{Nap. log } (10^7 - 1) = 1$. This I admitted in my former reply.

COLORADO SPRINGS,
June, 1916.

II. RELATING TO THE QUADRATIC FACTORS OF A POLYNOMIAL.

BY O. E. GLENN, University of Pennsylvania.

The process of finding by trial the rational roots of an equation $f(x) = 0$ with integral coefficients, depending upon the resolution of the last coefficient into its prime factors s_1, s_2, \dots and the first coefficient into its prime factors r_1, r_2, \dots , and testing by division the possible factors $x - s/r$, is given as an isolated method in books on elementary theory of equations, with no suggestion that it is capable of extensions. As an incident in a published paper¹ on Degenerate Curves I have shown, however, that this process is a particular manifestation of a general one in which, by a finite number of arithmetical trials, one can determine any polynomial $g(x)$ with integral coefficients which is a factor of $f(x)$. As applied to quadratic factors of $f(x)$, and especially when $f(x)$ is a quartic, the extended method is very useful.

By multiplying the roots of $f(x) = 0$ by a properly chosen integer we can assume it in the form

$$f(x) = x^n + p_1x^{n-1} + \dots + p_n = 0,$$

where the p 's are integers. Let a quadratic factor of $f(x)$, with integral coefficients, be $x^2 - \xi x - \eta$. If the roots of $f(x) = 0$ are $-x_1, -x_2, \dots, -x_n$, we may assume $-\xi = x_1 + x_2$.

¹ *Amer. Journal of Math.*, vol. 32 (1910), p. 79.

We have

$$P_4 = -2^3.83, \quad p_4 = -2.3,$$

and 6 is a superior limit to the roots. Hence

$$\xi = \pm 1, \pm 2, \pm 4, \pm 8; \quad \eta = \pm 1, \pm 2, \pm 3, \pm 6.$$

But $\xi_1 + \xi_2 = 6$, hence the pair (ξ_1, ξ_2) is either $(-2, 8)$ or $(2, 4)$, *i. e.*, $\xi = 2$ or -2 . The maximum number of unsuccessful trials is 15. With $\xi = 2$, $\eta = 6$, we have, employing an obvious extension of synthetic division,

$$\begin{array}{r} 1 - 6 + 3 + 22 - 6 \mid 2 + 6 \\ \underline{1} \quad \underline{2} \quad \underline{6} \\ -4 - 8 - 24 \\ \underline{1} \quad \underline{2} \quad \underline{6} \\ 0 \quad 0 \end{array}$$

Thus the remainder $ax + b$ is zero and

$$f(x) = (x^2 - 2x - 6)(x^2 - 4x + 1).$$

Further illustrations of quartics with irrational roots follow.

Let

$$f(x) = x^4 - x^3 - 6x^2 + 5x + 3 = 0.$$

Then

$$P_4 = 2, \quad p_4 = 3; \quad \xi = \pm 1, \pm 2; \quad \eta = \pm 1, \pm 3.$$

The only pair (ξ_1, ξ_2) for which $\xi_1 + \xi_2 = 1$ is $(-1, 2)$. Hence the maximum number of unsuccessful trials is three. We find

$$f(x) = (x^2 + x - 3)(x^2 - 2x - 1).$$

Exercise 1.—Show that the equation

$$x^4 + 4x^3 - 4x^2 - 17x + 10 = 0$$

can be resolved by less than nine trial divisions, and that the roots are $\frac{1}{2}(-1 \pm \sqrt{21})$, $\frac{1}{2}(-3 \pm \sqrt{17})$.

Exercise 2.—Solve the equation

$$x^4 + 4x^3 + 4x^2 - 16x + 28 = 0.$$

NOTES AND NEWS.

SEND ALL COMMUNICATIONS TO D. A. ROTHROCK, Indiana University.

The Secretary of the ASSOCIATION, Professor W. D. CAIRNS, is on leave of absence from Oberlin College during the present academic year. Until further notice he will be in residence at the University of Chicago and his official address will be 5465 Greenwood Avenue, Chicago, Illinois.